



**Postgraduate Day
April 20th**

BULLETIN

of the
**Mahoning
County
Medical
Society**

Vol. IX
April

No. 4
1939

FEED THE BABY

Dilution makes cow's milk more digestible — by attenuating somewhat the tough curd which characterizes cow's milk during digestion. But the milk-water-sugar formula still forms an appreciable curd . . . and the nutritional value of the mixture is further reduced by the dilution. Thus the usual "modification" as made in the home, if passably digestible, is commonly too low in caloric value. (Or too high in carbohydrate). ¶ Similac is cow's milk modified by laboratory methods. Not only are the fats, proteins and carbohydrate adjusted to approximately the percentages and nutritional values as in breast milk, but in addition the salt balance of the cow's milk is altered . . . SO THAT SIMILAC, LIKE BREAST MILK, STAYS LIQUID DURING DIGESTION. This is true of Similac no matter what concentration of the milk nutrients you find it advisable to prescribe. Because of this, the use of Similac reduces the occurrence of the difficult feeding cases which frequently result from underfeeding.

SIMILAC



M & R DIETETIC LABORATORIES, INC., COLUMBUS, OHIO

Similac is not advertised to the laity and no directions appear on or in the trade package.

Similac is made from fresh skim milk (casein modified) with added lactose, salts, milk fat, and vegetable and cod-liver oils.

COMMITTEE CHAIRMEN—1939

Name	Committee	Phone	Address
H. E. PATRICK	Editorial	4-2661	138 Lincoln Avenue
JOHN N. McCANN	Program	4-2772	2724 Mahoning Avenue
WALTER K. STEWART	Public Relations & Economics	4-1974	Stambaugh Building
E. C. RINEHART	Sub-Com. on Economics	5-2215	130 Bridge St., Struthers, O.
W. H. EVANS	Sub-Com. on Public Relations	4-2147	Dollar Bank Bldg.
JOS. C. HALL	Sub-Com. on Indigent Relief	6-6568	243 Lincoln Avenue
MORRIS ROSENBLUM	Publicity	3-9113	Home Savings & Loan Bldg.
C. A. GUSTAFSON	Correspondent	7-7224	101 Lincoln Avenue
D. M. ROTHROCK	Membership and Attendance	3-5688	1509 Market Street
DEAN NESBIT	Social	6-6703	Home Savings & Loan Bldg.
LEWIS K. REED	Lay Education	3-2422	1920 Market Street
GEO. M. McKELVEY	Housing & Library	7-7224	101 Lincoln Avenue
D. H. SMELTZER	Medico-Legal	6-6568	243 Lincoln Avenue
EDWARD J. REILLY	Postgraduate Day	3-0631	249½ Robinson Rd., Campbell
O. J. WALKER	Legislative	7-6176	Home Savings & Loan Bldg.
GORDON G. NELSON	Public Health	6-6844	138 Lincoln Avenue
LAWRENCE SEGAL	Auditing	6-6743	701 Himrod Avenue
WM. D. COLLIER	Field Army Against Cancer	4-4521	St. Elizabeth's Hospital

Vol. IX—No. 4

CONTENTS

April, 1939

COMMITTEE CHAIRMEN	97
ADVERTISERS' LIST	98
MEDICAL CALENDAR	99
PRESIDENT'S PAGE	101
RECENT ADVANCES IN THE SURGICAL TREAT- MENT OF PULMONARY TUBERCULOSIS	103-111
YOUNGSTOWN SYMPHONY SERIES	111
POSTGRADUATE DAY— FACULTY AND PROGRAM	112-113
MEDICAL CRIER	115
SECRETARY'S REPORT	115-117
MEDICAL MANAGEMENT OF PEPTIC ULCER	119-127
NEWS	128

ADVERTISERS' LIST

Patronize them—they make the
"Bulletin" possible

Baker Laboratories	116
Blair's Dry Cleaning.....	116
Buck, Maynard A.	124
Central Square Garage.....	108
Cross Drug Stores.....	118
Defiance Milk Products Co.	116
Denver Chemical Mfg. Co.	122
Dobson's	128
Endo Products, Inc.	120
Fair Oaks Villa.....	100
Foster, Helen Mantle.....	120
Giering's	128
Golden Age Gingerale.....	120
Heberding's	108
Hockstadt, R. L.	106
Isaly's	102
James & Weaver.....	124
Jones Metabolism Equip. Co.	106
Jones Pharmacy	120
Lyons Physician Supply Co.	110
Massachusetts Indemnity Ins. Co. ..	122
McKelvey's	98
Mead Johnson & Co.	Cover
Mercer Sanitarium	118
Merrell, Wm. S., Co.	Cover
Printz Co.	114
Renner's	122
Schmidt, Paul	120
Scott Company	104
Similac	Cover
Strouss-Hirshberg's	126
Thornton's	126
Treudley, H. H., & Co.	120
White Drug Co.	104
Wyeth & Bro., John.....	108
Youngstown Arc Engraving Co.	128
Youngstown Sanitary Milk.....	124
Zemmer Co.	126

ARROW



SPARE ROOM— Modern Conveniences!

MEN who wear Arrow shorts say they're the most comfortable they ever wore. Arrow shorts have *more room* . . . have *no center seam* to chafe your crotch . . . and they're *Sanforized*, the fabric is 99¾% shrink-proof! Come in and get some today.
65c, up.

See Our Complete
Line of
Arrow Ties
and Shirts
in the
Street Floor

MEN'S STORE

McKELVEY'S

ARROW

BULLETIN *of the*

Mahoning County Medical Society

A P R I L

1 9 3 9

Published monthly at
787 Wick Avenue, Youngstown, Ohio.

Annual Subscription, \$2.00.

WM. M. SKIPP, M. D., President

JOHN NOLL, M. D., Secretary

R. B. POLING, M. D., President-Elect

ELMER H. NAGEL, M. D., Treasurer

H. E. PATRICK, M. D., Editor

MEDICAL CALENDAR

April 20—Postgraduate Day—U. of P. group, Drs. Wolferth, Pendegrast, Stokes, and Kern.

May 16—Intern Competition, Youngstown Club.

June 20—Speaker, Dr. August A. Werner, Endocrinology.
Youngstown Club, 8:30 P. M.

July 20—Golf Party, So. Hills Country Club.

August 17—Golf Party, So. Hills Country Club.

September 14—Corn Roast and Clam Bake, Bert Millikin's Farm.

September 19—Speaker, Dr. Walter M. Simpson, Artificial Fever Therapy.
Youngstown Club, 8:30 P. M.

FALL LECTURES

October 2, 3, and 4—By Dr. Bernard Fantus.

Monday—11:00 a. m. to 1:00 p. m.: Prescribing of Pleasant Medication.

Tuesday—11:00 a. m. to 1:00 p. m.: Therapeutic Use of Fluids.
8:30 to 9:30 p. m.: The Therapy of Insomnia.

Wednesday—11:00 a. m. to 1:00 p. m.: The Therapy of Colon Stasis.
8:30 to 9:30 p. m.: The Therapy of Cough.

RADIO TALKS

April 7—What's Behind a Prescription? - - - Dr. Samuel Schwebel

April 14—What Is Occupational Therapy? - - - - Dr. L. S. Shensa

April 21—Tired Legs from Altered Circulation - - Dr. M. H. Steinberg

April 28—Medical Ethics - - - - - Dr. Morris Belinky

May 5—Cosmetics and Common Sense - - - - Dr. Samuel Zoss

May 12—Invisible Handicap - - - - - Dr. Stanley Myers

May 19—Prevention of Blindness - - - - - Dr. William F. Hatcher

A Modern Institution for the Care of Nervous and Mental Patients Alcohol and Drug Addiction

A majority of the patients are referred to Fair Oaks Villa by members of the medical profession and it is only through the journals of the profession that any publicity is sought. The interests of FAIR OAKS VILLA, as an Institution, are completely dominated by those who have high regard and respect for the tenets of professional ethics.

The homelike atmosphere of Fair Oaks Villa is afforded by the pleasant rooms, by the eight and one-half acres of attractive lawns, picturesque landscaping and restful surroundings.



G. D. TIPTON, M. D.
Medical Director

A. B. BOSMA, M. D.
Assistant Physician

R. E. PINKERTON, M. D.
Consultant

E. D. McGOWAN, President

E. A. PAISLEY, Business Mgr.

Fair Oaks Sanitarium is a member of the American Hospital Association and the Central Neuropsychiatric Hospital Association.

PRESIDENT'S PAGE

Postgraduate Day

The biggest show of the year will be staged in Youngstown April 20, 1939. This will be the Twelfth Annual Postgraduate Assembly of the Mahoning County Society and from all indications it will be the greatest day the Society has ever sponsored.

May I extend to our guest speakers, Dr. Wolferth, Dr. Pendergrass, Dr. Kern, and Dr. Stokes, of the University of Pennsylvania, the Society's heartiest and sincere greetings. May they find us an attentive audience and appreciative of their efforts put forth for our benefit.

The Mahoning County Medical Society extends a cordial welcome to all members of the Profession near and far, and especially those of Ohio and Western Pennsylvania. We can without reservation promise a full day of valuable education and real enjoyment, meeting old and making new friends. The program is excellent, the food, none better, in fact the day's entertainment is complete in every detail. The Postgraduate Committee with the aid of the Program, Entertainment and Publicity Committees, will assure no hitch in the proceedings from beginning to end.

These Committees function so well there is no need for a directing head or a moment's worry, for the show is bound to be a success. They are all veterans, have been well trained, and no amount of effort on their part will be lacking so the show will go on full of vim and vigor, never a dull moment.

We solicit helpful criticism from our members and guests. Do not hesitate to call to the attention of any member of the committees or the officers of the Society any function that has not been to your liking.

It is our aim, as in the past, to have this the best meeting of its kind in this section of the country. We will have visiting doctors, new faces, whom we will not know. It is therefore the duty of each and everyone of us to extend a friendly hand of fellowship, help him to know our Society, its members, our city, and tell him of our aims and what our Society stands for in teaching scientific medicine and its ethical practice.

We must not forget our *Bulletin*, the mouthpiece of our Society, that through its ever-present reminding, our Postgraduate Day is made possible. Our Editor and co-workers never cease to be at work on the duties of our Society.

To our advertisers and exhibitors we extend a hearty welcome. Their displays are as good if not better than many State and National meetings. The business firms represented are honest, ethical and conscientious representatives of the trades that are essential to the practice of medicine.

The Mahoning County Medical Society sends greetings and a cordial invitation for all to attend this Twelfth Postgraduate Day Assembly, April 20, 1939, which will be held at the Ohio Hotel, Youngstown, Ohio.

May we see you there? To the members of our Society, this is your Postgraduate Day. It cannot be a success and function without your whole-hearted support. The Committees can arrange the meeting but you must attend if victory would be ours.

— WM. M. SKIPP, M. D.

Cheplin's "The Original"
B. ACIDOPHILUS MILK

Accepted by the Council of the
American Medical Association



Distributed by
The
Isaly Dairy Co.

The first commercial culture of *Bacillus Acidophilus*, accepted by the Council of the American Medical Association, was so-called *Acidophilus Milk*. Because of its priority in recognition by the medical profession, *Acidophilus Milk* has been set up as the standard, in terms of which other commercial *Acidophilus* cultures are evaluated. *Acidophilus Milk* is recommended in the treatment of chronic constipation, colitis, dysentery, fermentative diarrhea, and intestinal putrefaction. In general, the *Acidophilus* treatment may be employed in correcting disorders arising from the absorption of poisons originating in the human digestive tract.

Through a special daily distribution system maintained by the Cheplin B. *Acidophilus Milk* Manufacturers, any orders placed through The Isaly Dairy Company (please use phone 6-5246) will receive prompt attention.

ISALY'S
DAIRY SPECIALISTS

RECENT ADVANCES IN THE SURGICAL TREATMENT OF PULMONARY TUBERCULOSIS

JOS. KEOGH, M. D.

The management of pulmonary tuberculosis is without question the most important problem with which thoracic surgery is confronted today. When one considers that there is an average of seventy deaths per hundred thousand inhabitants and seven tuberculosis patients for every death, we can readily see that this is true. While lung abscess, carcinoma of the lung, tumor of the lung and mediastinum or bronchiectasis, all present more spectacular results, figures just quoted make these problems of secondary importance.

The treatment of pulmonary tuberculosis has undergone radical changes during the past few years. From being ultra-conservatively symptomatic, expectant, climatic and sanitorial, the pendulum has swung to the other extreme in many places and is entirely surgical. It is the purpose of this paper to present an attempt at a more rational balance between the two extremes and to emphasize the indications for surgical intervention. It would be impossible in such a short paper to give any detailed technical description of each of the procedures used but there will be space I believe to indicate the method most suitable for use in the different types of cases.

As has been pointed out by the late Dr. Coryllos, three major morphological changes may occur in the parenchyma of the lung infected by the tubercle bacillus:

1. Infiltration and edema (allergic phase).
2. Caseous necrosis of the affected lung parenchyma (destructive phase).
3. Development of fibrous tissue in the involved areas (repairative stage).

Caseous necrosis of a portion of the parenchyma leads to the formation of tuberculous cavities following the elimination of the caseated material through the bronchi. These cavities,

which are holes in the lung containing air, represent the most characteristic features of pulmonary tuberculosis and it is toward the closure of these cavities that surgical management is directed.

A study of the mechanism whereby closure of cavities is effected has resulted in many theories. Such answers as, "Nature has a tendency to produce closure of cavities," cannot be accepted. Some men have advocated the presence of a double system of concentric and radiating fibres, one of which tends to produce retraction and closure and the other, the opposite effect. Unfortunately, these fibres have never been visualized under the microscope. Others have based their theories on the changes in elasticity of the lung. For these, the elasticity is increased while for still others, it is decreased and even yet further, for Bendove, the elasticity of retraction is increased whereas, the elasticity of expansion is decreased. Since the elasticity of the lung is a function dependent upon the elastic fibres of the lung and since destruction of elastic tissue takes place in all diseases of the lung in general and in tuberculosis particularly, it would seem that the theory was defeated before it started. However, when one considers the physiology of the lung and compares it with the pathological physiology of the lung, the answer becomes obvious. The late Dr. Coryllos, in several recent papers, has advanced proof of the simplicity of this mechanism. A cavity appears spherical in the living subject by virtue of the atmosphere of pressure within, since the cavity is in direct communication with an open bronchus, being greater than that exerted by the surrounding lung. The pressure exerted by the surrounding lung is, of course, only equal to the negative intrapleural pressure. Closure of



C. W. Ourant

C. W. OURANT

TRUSS FITTER

is now showing the new

Men's Wide-Band Supporters

—those helpful, healthful, cool elastic belts that also add to the personal appearance.

Postgraduate Day he will display those new thin elastic hose for men and women by Bauer & Black.

WHITE'S DRUG STORES

Dependable Prescription Druggists

POSTGRADUATE DAY — APRIL 20

See these Smart, Wide Brim
Knox Hats for Spring

You'll like their mannish look . . .

their rich quality . . . and

they're now available at \$5.00.

Other grades \$7.50 and \$10.

The Scott Company

32 North Phelps Street

the communicating bronchus results in gradual absorption of air, collapse of the cavity and thus closure of the cavity. The changes in gaseous contents of the cavity before, during and after bronchial closure have been studied as well as measurements taken of the gaseous pressure within the cavity. All results obtained from this study indicate that closure of the cavity is effected by closure of the draining bronchus.

This discussion leads up to the consideration of the indications for collapse therapy in pulmonary tuberculosis because all collapse therapy is directed toward the closure of cavities. In the following discussion, the classifications presented by Ulmar, Ditter and Ornstein will be used. The acute exudative or allergic phase still belongs and should be managed along established conservative lines. Radical bed rest, good food, hygienic measures as well as general supportive measures remain the treatment of choice in this group. Many of these lesions, given time, will clear of themselves. Even aside from this, any collapse measures attempted on the acute exudative lesion may not only lead to the development of cavernous lesions but also may result in actual ulceration through and into the pleura or, finally, to a bronchogenic or hematogenous spread. It is our opinion that all such lesions should be carefully watched for at least a period of six months. During this length of time, the involved area will usually either show signs of regression or will go on to a definite caseous pneumonic type of disease.

Chronic productive pulmonary tuberculosis (Acinous Nodosa Form of Aschoff) presents a similar problem. Surgery on this type of lesion will produce spectacular results but actually most of these lesions will clear of themselves given adequate opportunity. Cases which have a caseous pneumonic lesion on one side often have productive changes on the contra-lateral or opposite side which

automatically clear once the cavity has been closed.

Finally, we must consider the caseous pneumonic or chronic ulcerative form of pulmonary tuberculosis. By that we mean, those cases with open cavities and positive sputum. Even in this group, immediate surgical intervention is actually extremely radical but conversely as well such conservatism becomes even more radical when allowed to go on too long. About ten to twelve percent of caseous pneumonic lesions close spontaneously. In the face of no spread of disease, no progressive increase in the size of the cavity, they should be given a reasonable length of time to effect a closure. However, unnecessary delay is the most dangerous treatment one can offer a patient. The possibilities of such a thin walled cavity becoming a thick, giant or tension cavity (cavity over 8 cm. in diameter), source of bronchogenic spread to other lobes on the same side or contra-lateral side are far too great.

The indications for surgery narrow themselves for all intentions and purposes to the treatment of cavitation in the lung, hemorrhage and empyemata. The various surgical procedures utilized depend almost entirely upon the type and distribution of such lesions. This, of course, excludes from consideration that group having either a toxic tuberculous empyema, a mixed infection tuberculosis and pyogenic empyema with or without a broncho-pleural fistula. Briefly, all of this latter group requires immediate surgical intervention of a most radical nature.

As mentioned above, the choice of the procedure depends entirely upon the type and distribution of the lesion. Procedures now in vogue are:


1. Pneumothorax.
2. Closed pneumonolysis. Closed pneumonolysis, an operation to complete a pneumothorax by severing the adhesions suspending the lung to the

For Accuracy and Simplicity in making
Metabolism Tests install the
**NEW JONES
MOTOR-BASAL UNIT**

Thousands of users all over the
country enthusiastically favor
the **NEW OUTSTANDING
FEATURES** of this model

- ✓ SIMPLIFIED TECHNIQUE AND
NO CALCULATIONS
- ✓ ECONOMY OF OPERATION
- ✓ LIFE-TIME GUARANTEE
- ✓ WATERLESS
- ✓ NOISELESS, ALL-ELECTRIC,
BALLBEARING EFFICIENCY

Mail in coupon below and take
advantage of our offer of a **FREE
DEMONSTRATION** of its exclusive
features and a trial period of the
apparatus without any cost or obli-
gation. This service is coupled with
instructions that will greatly enlarge
the scope of the use of the B.M.R.
in every general examination.



Accepted by the
Council on Physical
Therapy of the
A. M. A.

Small Monthly
Payments Out
of Income Will
Enable You to
Own It At Once

Jones Metabolism Equipment Co., 1870 E. 90th St., Cleveland, O.
Gentlemen: Please send me
 Booklet on "Accurate Basal Metabolism Testing and Clinical
Applications"
 Full details on your 7-day trial offer
NAME _____
ADDRESS _____
CITY AND STATE _____

POSTGRADUATE DAY — APRIL 20

R. L. Hockstad

PRESCRIPTION SPECIALISTS

268 Lincoln Avenue

Youngstown, Ohio

Phones 40107 and 40108

Office Supplies Reasonably Priced

100% Ethical

•

No Counter Prescribing
No Substitution
No Therapeutic Discussions
Nothing but Prescriptions

•

Our R. Price is always fair.

No Extra Charge for delivery if your R. is phoned to us.

chest wall. Actually, while the operation is apparently very simple, its performance requires more anatomical knowledge, keener judgment and better technique than do any of the apparently more radical procedures. Even the smallest error in judgment may result fatally through the development of a tension pneumothorax or a mixed infection empyema.

3. Phrenic nerve interruption, either temporary by crushing or permanent by section.

4. Apicolysis with packing or plombage.

5. Extra-pleural pneumothorax.

6. Thorocoplasty.

7. Thorocoplasty combined with Schedde for the obliteration of a chronic empyema space.

8. Lobectomy and pneumonec-tomy.

Rather than consider each of these procedures separately, it would seem better to consider the pathology requiring therapy and simply to name the procedure best suited for its control. Unilateral apical cavities are best controlled by apical thorocoplasty either in one or two stages. While this may seem extremely radical, actually, it is quite conservative. Pneumothorax therapy on such a lesion must be continued for the usual length of time. The most accepted time is three to five years. During this time, the patient is continually subject to the dangers of a spontaneous pneumothorax, which may be either traumatic or ulcerative. They are also subject to the development of a tuberculous empyema or even worse, a mixed infection empyema with bronch-pleural fistula. Thorocoplasty operations, at the present time, are not as deforming nor is the mortality rate as high as in previous years. When one considers that better than eighty percent of such unilateral lesions can be controlled by this procedure, meaning that within a period of six to eight weeks the cavity is usually closed and within a

six-month period, the patient can be safely discharged. Then that which appears to be radical becomes conservative from the patient's standpoint, the public health standpoint and the cost of hospitalization for such an individual. Pneumothorax, on the other hand, should first be tried on cases demonstrating lesions disseminated throughout one lung. In the event that only a partial collapse can be obtained, this may be completed by means of a closed pneumolysis. It is often surprising how much can be done to complete what may appear to be a very inadequate pneumothorax by severing multiple adhesions in this manner. In all cases, where pneumothorax fails and cannot be completed by intra-pleural pneumolysis, thorocoplasty becomes the procedure of choice. This is usually a complete thorocoplasty requiring three stages.

Bilateral pulmonary tuberculosis, caseous pneumonic form, may be controlled in one of several ways. Bilateral apical lesions are best controlled by bilateral apical thorocoplasty.

In the presence of an apical lesion on one side and multiple cavities on the contra-lateral side, pneumothorax may be attempted on the side with the apical lesion and then, thorocoplasty done in the presence of the pneumothorax on the contra-lateral side. These cases must be handled with great care because of the possibility of a traumatic or actual spontaneous pneumothorax which may occur during the course of refills following operation. Bilateral, equally disseminated lesions may be controlled by bilateral pneumothorax which in many instances must be completed by unilateral or bilateral closed pneumolysis. Another recent adjunct to bilateral collapse therapy has been the introduction of extra-pleural pneumothorax. There are certain cases with bilateral lesions where pneumothorax is impossible on both sides and where bilateral thorocoplasty would be beyond the limit



LET ME HAVE IT!

Doc Says—

There is just one way
to keep my batting aver-
age up

drink

MILK and CREAM

from

HEBERDING'S INDIAN CREEK FARM MILK

Florence L. Heberding

Phone 2-2344

POSTGRADUATE DAY — APRIL 20

Amphojel
Kaomagma
Ergoklonin
Citri Cerose
Wyamoids
Elixir Bewon
Silver Picrate - Supp.
" " Pow.
Mucara

JOHN WYETH & BRO.

N. G. Yates

Representative

3057 W. BAILEY ROAD
CUYAHOGA FALLS, OHIO

**• PROMPT AND
COURTEOUS SERVICE**

•
While Your Car Is Parked

Have It

Washed and Lubricated

•
Central Square Garage

Wick and Commerce Sts.

Phone 3-5167

of that person's vital capacity. These cases were, therefore, outside the realm of thoracic surgery. With this new operation, following the resection of a short length of one rib, a pneumothorax space is operatively created by apicolysis in the extra-fascial plane. After an air-tight closure, this space is then filled with air and periodical refills given as in the management of an ordinary intra-pleural pneumothorax. Ordinarily, the first refills are quite difficult but once the space is established, the pneumothorax is easily maintained. Positive pressures are used.

The phrenic nerve operation has enjoyed a vogue over many years but in recent years is being abandoned by even those who formerly most strongly advocated its use. In our group of cases, we found one type of lesion most susceptible to control by this operation. These lesions are mid-field cavities on which a pneumothorax has had appreciably no effect. By combining the phrenic nerve operation with the pneumothorax, a closure has been effected. Elevation of the diaphragm is also of aid in many instances to help decrease the size of an empyema space.

Although, no details of technique can be gone into in a paper of this length, it seems only just that a few of the more recent changes in thorocoplasty technique should be mentioned. Instead of the old Sauerbrauch operation with the resection of short lengths of many ribs, we now do multiple stages with the resection of much greater lengths of rib. Complete thorocoplasty for a lung equally infected throughout, usually demands three stages. The first stage requires the sub-periosteal resection of total lengths of the first three ribs. During the second stage, three and sometimes, four ribs are removed while the third stage is limited to a maximum of three ribs because of the marked paradoxical respiration which follows removal of more than three. A new modification for the

control of giant apical cavities has recently been introduced by Dr. Carl Semb. Briefly, the operation is done as follows: Following the resection of the first three ribs, the apex is decollated by blunt dissection to the level of the fourth rib posteriorly. A few of the surgeons using this procedure fill the dead space thus created with saline to insure maintenance of the drop. In our experience, this dead space fills completely within six hours post-operative with a sero-sanguinous type of fluid. Up to the introduction of this modification, thorocoplasties on giant cavities had been almost a total failure and it is in this group, that most of the radical revisions had to be done. The Semb modification carries with it a slight increase in mortality but when one considers that formerly none of these cavities were closed without radical revisions, which in themselves carry a mortality of thirty-two percent, it seems the additional risk is well worth while. The most recent technical addition is one devised by the late Dr. Coryllos. Thorocoplasties on children under sixteen years of age had been more or less abandoned because of the marked scoliosis. The resulting deformity was often more severe than the disease. By combining spinal fusion and thorocoplasty done at the time of thorocoplasty, this complication has been overcome.

The other procedures such as apicolysis with plombage, oleothorax, etc., have all enjoyed a brief popular term but are not being generally used at the present time. Lobectomy and pneumonectomy have been unsuccessful up to very recently. Some few cases have been reported as successful but they are far too few to draw any conclusions as yet. Up to now, the complications following these procedures have been too serious to warrant their use.

There is no question that collapse therapy has markedly influenced the mortality rate in pulmonary tuberculosis. Spontaneous closure of ca-

WELCOME POST GRADUATES

VISIT OUR DISPLAY

WE SPECIALIZE IN:

SHORTWAVE
DIATHERMY

ELECTRIC
CENTRIFUGES

COLD QUARTZ
LAMPS

INFRA RED
LAMPS

NATIONAL CARBON
LAMPS

ALLISON OFFICE
FURNITURE

SUCTION
MACHINES

SURGICAL
INSTRUMENTS

TRUSSES
BELTS and ELASTIC
HOSIERY

ANALGESIC
TABLETS

PHENALGESIC
TABLETS

PHENOCOLATE
TABLETS

HYPERTENSION
TABLETS

ELIXIR OXGALL
and PANCREATIN

RECONSTRUCTIVE TONIC
with VITAMIN B₁

ELIXIR PHENOBARBITAL
DOUBLE STRENGTH
RED or GREEN

IODYL
Effective Throat Swab

ALL STANDARD
MANUFACTURES
SPECIALTIES

LYONS PHYSICIAN SUPPLY CO.

26 Fifth Avenue

Youngstown, Ohio

Phone 4-0131

vities takes place in about ten to twelve percent of all cases. About twenty percent of all cases receiving pneumo-thorax therapy becomes negative. Closed pneumonolysis completes about seventy percent of unilateral cases on which it is tried and approximately twenty-five percent of bilateral bases. Sixty-two percent of these people develop an effusion. Two and one-half percent develop a tuberculous empyema and about two percent a mixed infection empyema. In 529 cases or 1275 operations of thorocoplasty at Sea View Hospital, New York, the general mortality was 18.3% per patient mortality or 7.6% for operation. This, however, includes thirty-two deaths associated with operation for tuberculous empyema, mixed infection empyema and revision. The actual mortality for straight thorocoplasty is therefore, operatively 5.1%. This includes both unilateral and bilateral cases. The operative mortality for unilateral cases alone is 3.3%. Of this group, exclusive of the empyemas, 82% had conversion of sputum from positive to negative.

In conclusion, I should like to emphasize that of all those actively engaged in the control of pulmonary tuberculosis, the thoracic surgeon is the first to realize that surgery is not the final answer to the solution of the problem. Even further than that, it is not by any manner or means a direct attack upon the disease as a public health problem. The ultimate cure when it comes will be medical. However, collapse therapy in its various forms is, to date, the most effective means of controlling the individual case and in well regulated groups with close coöperation between the roentgenologist, the medical internist, the ear, nose and throat group and the surgeon, there has been and increasingly is found, a definite decrease in mortality. This is accompanied by a decrease in hospitalization and an even more definite and more important increase in the number of patients treated and in the number discharged under conditions suitable for their return to society with safety.

YOUNGSTOWN SYMPHONY ANNOUNCES 1939-1940 SOLOISTS

Soloists of unusual interest have been engaged to appear with the orchestra in addition to the local soloists Carmine Ficocelli, Jacob Huebert and Herman Gruss, who perform Beethoven's Triple Concerto on the February concert, five world famous solo attractions have been engaged. The list includes The Don Cossack Male Chorus, Serge Jaroff, conductor, who appear on the gala opening concert in an all-Russian program; Silvio and Isabel Scionti, brilliant duo-pianists who will play a two piano concerto with the orchestra; Moriz Rosenthal, greatest of Franz Liszt's pupils, who will be on his Golden Jubilee Tour with his famous gold piano; Helen Jepson, glamorous young American soprano

of the Metropolitan Opera and native of Columbiana and Akron, Ohio. Albert Spalding, America's foremost violinist is the soloist on the closing concert in April.

Mr. Spalding appeared with the orchestra two years ago and many requests from patrons for a re-appearance prompted the engagement for next season.

The same low season prices will be in effect next year and subscriptions are now being received at the Symphony headquarters—307 Union National Bank Bldg., in Youngstown. The 1939-1940 series will again be sponsored by the Junior Chamber of Commerce.

POSTGRADUATE DAY FACULTY



Dr. Charles C. Wolfert

Professor of Clinical Medicine, School of Medicine, University of Pennsylvania.
 Administrator, Robinette Foundation.
 Member of Medical Staff, Hospital of the University of Pennsylvania.
 Chief of Cardiac Section, Medical Division, University Hospital.
 Consulting Physician, Fitzgerald Mercy Hospital
 Consulting Physician, Underwood Hospital
 Consulting Cardiologist, Jewish Hospital.
 Consulting Cardiologist, St. Joseph's Hospital.

Director of the Department of Radiology, University of Pennsylvania Hospital.

Professor of Radiology in the Graduate and Undergraduate Schools of the University of Pennsylvania.



Dr. E. P. Pendergrass



Dr. Richard A. Kern

Graduate of the University of Pennsylvania—1914.

Professor of Clinical Medicine.

Head of the Allergy Clinic, University Hospital.

Haverford College, A. B., 1916.
 School of Medicine, University of Pennsylvania—1920.

Pediatric Staff—Physician-in-Chief—Children's Hospital.

Society for Pediatric Research—One of the founders and former president American Pediatric Society.

American Academy of Pediatrics, Co-Chairman for Pennsylvania.



Dr. Joseph Stokes, Jr.

A Program You Cannot Afford to Miss!

By

GROUP FROM UNIVERSITY OF PENNSYLVANIA

Twelfth Annual

Postgraduate Assembly

APRIL 20, 1939



MORNING SESSION

- 9 to 10 A. M. Cardinal Features of Coronary Disease - - - - - Dr. C. C. Wolferth
- 10 to 11 A. M. Recent Advances in the Study and Treatment of Certain Virus Diseases - - Dr. Joseph Stokes
- 11 to 12 A. M. General Principles of Allergy and Clinical Manifestations of Allergy (Part 1) - - Dr. R. A. Kern

LUNCHEON

AFTERNOON SESSION

- 2 to 3 P. M. Silicosis and Silicosis with Infection - - - - - Dr. E. P. Pendergrass
- 3 to 4 P. M. Clinical Manifestations of Allergy and Principles of Diagnosis and Treatment— (Part 2) - - - - - Dr. R. A. Kern
- 4 to 5 P. M. Prophylaxis and Treatment in Certain Acute Infectious Diseases - - - Dr. Joseph Stokes

EVENING SESSION

- 6:30 P. M. Dinner
1. Roentgen Therapy of Infection - - - - - Dr. E. P. Pendergrass
 2. Physical Examination of the Heart with Special Reference to Heart Sounds - Dr. C. C. Wolferth



"Printztonian" Suits

Here are suits designed to give plenty of hard service. Fine worsteds of long-staple yarns expertly tailored will take all the abuse you give them . . . and come back for more . . . Get the added service *Printztonians* afford!

Talon Closure on Both Trousers!

\$35⁰⁰
Two
Trousers

The MODERN
PRINTZ'S



THE MEDICAL CRIER

A Page of Sidelights, News and Views in the Medical Field

● "One-third of the population is unable to obtain medical service." Where do they get that stuff? What factual study, what widespread survey has brought to light the existence of this deplorable state of affairs? Has this awful pronouncement so frequently used by proponents of Socialized Medicine any basis in fact? Certainly not around here. Every member of this Society is certain of that. But maybe we don't know about conditions in other parts of the country. Maybe the deep South or the far West are different.

Reports received on the American Medical Association Study of Need and Supply of Medical Care already cover a far larger sample of the population than has been covered by any previous survey. Questionnaires have been received from hospitals, nurses, health departments, schools, welfare and relief agencies, colleges and universities, fraternal organizations and pharmacists. Only one of the nine forms used in this survey has been directed to physicians and dentists. Not one significant source of first hand, reliable information as to medical conditions has been omitted, and so far there has been no sharp disagreement in the reports.

There is no locality in which any organization even suggests that one-third of the population is unable to obtain medical service.

The most general lack of medical care that is reported is with regard to insufficient appropriations for poor relief or governmental institutions. Isn't this significant?

● We saw with much pleasure the article on "Aesculapius" by Dr. Louis Deitchman reprinted with due credit to the M. C. M. S. Bulletin in the Westchester Medical Bulletin. It has been our privilege to print several of Dr. Deitchman's historical sketches, the most recent and instructive being

the story of Washington's last illness presented in case history form. A collector of old books and first editions, his ventures into medical history have always been interesting and authentic. Not to mention the witty doggerel which formerly appeared under the pen name of Theophrastus Bombastus. If properly urged, no doubt this scholarly fellow would give us more in similar vein.

● There seems to be a premium placed nowadays on smartness. The bright fellow, the keen wit and the wisecracker get the headlines. Jimmie Cagney, movie star in the current picture "Oklahoma Kid" says, "Listen, the strong take it away from the weak, but the smart take it away from the strong." So it goes, but in the long run character is greater than smartness, especially in our profession. A great business leader once said, "My son is not smart, but he has character. I can hire a hundred men who are smart, but a man with character is mighty nice to live with." The prestige of the medical profession is enhanced by the few who are brilliant, but its reputation and confidence are built upon the solid foundation of its general good character.

J. L. F.

SECRETARY'S REPORT

March, 1939

At the regular March Council meeting held on the 13th of the month, the members surveyed the new bookkeeping system and membership files as recently set up by Mary Herald, the assistant secretary. This system has been approved by W. C. Fisher, auditor, and will make for more efficient bookkeeping records.

A special Council meeting was called by President Skipp following the regular meeting on March 21, to nominate three doctors for each of the following boards:



The True Test of an Infant Food

is not the results it may produce during only a brief period—for the first ten days after birth, or perhaps for another month or two.

A good infant food will give good results continuously. That is why so many physicians prefer

BAKER'S MODIFIED MILK

Being a well balanced diet, with an adequate protein content, and fortified with all the essential vitamins (excepting C), Baker's MODIFIED MILK produces good results from birth until the infant is finally weaned from the bottle.

Advertised only to Physicians

Available in both powder
and liquid forms.



The Baker Laboratories — Cleveland

Other Baker Products

MELCOSE—a completely modified liquid milk. Very economical. A single can costs 15c & dilutes to a full quart.
MELODEX—Maltose and dextrin, an easily assimilated carbohydrate especially prepared to modify cow's milk. Price 49c.

POSTGRADUATE DAY — APRIL 20

EVAPORATED Morning Glory Milk



IMPROVES
COOKING
and
BAKING
"IT WHIPS"

Fine for Babies
and Invalids

**DEFIANCE
MILK PRODUCTS CO.**

RELY ON BLAIR'S WORKMANSHIP

Let the BLAIR driver take down your drapes, roll up your rugs, and move your upholstered furniture. They will be spotlessly cleaned and have that new appearance.

Call for estimates.

Phone 4-4227

Blair's

Responsible Cleaners and Furriers

(1) The Mahoning County Welfare Board;

(2) The Board of the Mahoning County Tuberculosis Sanitarium.

The following doctors were chosen: Dr. J. B. Birch, Dr. W. H. Evans, Dr. Barclay M. Brandmiller for the Welfare Board; Dr. C. R. Clark, Dr. L. S. Deitchman and Dr. E. J. Reilly for the Mahoning County Tuberculosis Sanitarium Board. These names were duly submitted to the Mahoning County Commissioners from which they will choose one for each board.

The regular monthly meeting for March was held on Tuesday evening, March 21, at the Youngstown Club.

The scientific program was presented by Dr. George M. Curtis, Chairman of the Department of Research Surgery at Ohio State University Medical College. He spoke on "The Iodine Metabolism in Thyroid Disease" and brought the work up to date citing interesting charts of Iodine Imbalance in Thyroid Disease. His work in this field is meticulous and shows very careful technique for accuracy. The presentation had many practical sides as well as of just scientific interest.

Following the scientific meeting a business meeting was held. The applications of the following doctors were read:

For Active Membership:

Dr. Wm. Peter Beckley
Dr. Milton Marvin Yarmy

For Associate, Class D:

Dr. John S. Goldcamp

Dr. Sedwitz moved that the Society approve the increase in State dues from \$5.00 to \$7.00, as explained in the January issue of the Ohio State Journal, and so instruct our State delegates to vote for the increase at the coming State meet-

ing. Motion was duly seconded and passed.

Dr. Tims moved that the Society approve the redistricting of the State, adding one new district, the Eleventh, as described in the January issue of the Ohio State Journal; and so instruct our State delegates to vote for the amending of the State Constitution so that this can be possible. Motion was duly seconded and passed.

Dr. Skipp then announced that the County Commissioners have requested the names of three doctors of the Mahoning County Medical Society from which one will be chosen for each of two boards, namely, the Welfare Board and the Board of the Mahoning County Tuberculosis Sanitarium. Dr. Tims moved that Council be named a committee to choose the six nominees. Motion was duly seconded and passed. Dr. Skipp then called for a Council meeting immediately following this regular meeting, because the names must be submitted to the Commissioners by the following day.

Dr. Reilly then appealed for publicity on Postgraduate Day.

Dr. Skipp discussed the suggestion of changing of half holiday from Thursday to Wednesday afternoons. There was no discussion.

Meeting adjourned.

Council has approved the applications for membership in the Mahoning County Medical Society of the following:

For Active Membership:

Dr. M. M. Szucs

For Class D Membership:

Dr. John Smith, Sebring, O.

Unless objection in writing to any of these applicants is filed with the Secretary within 15 days, they will become members of the Society.

DR. JOHN NOLL, *Secretary*.

THE MERCER SANITARIUM MERCER, PENNSYLVANIA

For Nervous and Mild Mental Disorders.
Located at Mercer, Pennsylvania, thirty miles
from Youngstown. Farm of one hundred
acres with registered, tuberculin-tested herd.
Re-educational measures emphasized, especially
arts and crafts and outdoor pursuits. Modern
laboratory facilities.

Address:

W. W. RICHARDSON, M. D., Medical Director
Formerly Chief Physician, State Hospital for Insane, Norristown, Pa.

POSTGRADUATE DAY — APRIL 20

Youngstown Owned and Operated

CROSS DRUGS

6 EAST FEDERAL STREET
In the Stambaugh Bldg.

1 WEST FEDERAL STREET
In Central Tower Bldg.



WE INVITE YOU TO VISIT OUR
PRESCRIPTION DEPARTMENTS

Five Pharmacists to Serve You.

Prescriptions Delivered to Your Patient.

THE MEDICAL MANAGEMENT OF PEPTIC ULCER*

L. S. SHENSA, M. D.

In this paper I shall attempt to review the various methods of treatment of peptic ulcer as practiced in several large clinics and also give an evaluation of their results.

The treatment of peptic ulcer requires a definite plan of education. Like the care of such diseases as diabetes, tuberculosis or chronic heart disease the full and intelligent co-operation of the patient must be gained if treatment is to be successful. The patient must not throw the whole burden of his disease on the physician's shoulders but he must cooperate intelligently with his doctor. We can never be sure that the peptic ulcer is permanently healed. Relaxation in therapy will result in an exacerbation of the disease in most instances. One of the most important points in the management of peptic ulcer is the modification of the patient's habits of living. We must secure tranquility of mind and body as far as his social and business conditions permit, nine hours of sleep at night, a mid-day rest of one-half hour and simple recreation during leisure hours are of vital and essential importance. A period of change away from the home environment is most valuable. Moderate work in the garden, music or some hobby will give a release from nervous tension. Frequently this treatment alone will produce relief from symptoms. Definite foci of infection should be eliminated at the proper time. Diseases of the pelvis or appendix may require surgical care.

Simple hypochromic anaemia will respond to adequate iron therapy, but it is best not to give iron by mouth until the painful stage of ulcer has passed.

The peptic ulcer patient should be sent to the hospital when he has

suffered from one or more severe hemorrhages, when x-ray examination shows that a penetrating ulcer or pyloric obstruction is not improved with due trial with ambulatory care, and if the pain is so severe that there is some question as to the correct diagnosis.

The ambulatory method of treatment of peptic ulcer may be employed if the patient of limited financial resources desires to be kept on the job. This reduces his anxiety about the welfare of his family. His general strength will be preserved by keeping him on an adequate diet from the very beginning of his treatment, and he will be spared a difficult period of adjustment to normal living which often occurs if he has spent a few weeks in bed or in the hospital. Finally, the best reason is that in this type of treatment, if intelligently carried out, the results are very good.

The indications for surgical intervention in peptic ulcer are quite definite:

1. In cases of intractable ulcer.
2. In those cases of gastric ulcer in which bi-weekly follow-up x-ray studies fail to show regression of the condition. All gastric ulcers are under suspicion of cancer until they show definite evidence of healing.
3. Pyloric obstruction which resists medical therapy.
4. Persistent or recurrence of massive gastric hemorrhage.
5. Acute or chronic perforating ulcers.

Medicinal Therapy:

It is seldom that unusual or complicated medicinal therapy is required. Despite some opinions to the contrary belladonna or atropine are the most satisfactory anti-spasmodics. As a rule pyloric spasm and the gastro-intestinal hypermotility will be definitely abated by the use of belladonna. At

*Read to the Staff of St. Elizabeth's Hospital at its regular monthly meeting on March 14, 1939.



**Endo
ESTROMONE**
ESTROGENIC
HORMONE
(OVARIAN FOLLICULAR)
Now in tablet form

**EFFECTIVE ORAL MEDICATION
IN OVARIAN THERAPY**

Endo Estromone, in tablet form, provides the same pharmacologic activity as Estromone in oil solution . . . permitting easy oral medication where hypodermic injection is undesirable, or as a supplement to intramuscular administrations.

Indications: Gonorrhoeal Vaginitis (in children), Senile Vaginitis, Functional Dysmenorrhoea (due to Estrin deficiency), Relief of Natural and Surgical Menopause, Involutional Melancholia, Breast Hyperplasia (lobular type), certain cases of frigidity.

Supplied in Tablets of 1,000 and 2,000. International Units, in packages of 20 and 100 Tablets.

ENDO PRODUCTS, INC.
395 Fourth Avenue New York, N. Y.

**FLOWERS FOR
ALL OCCASIONS**
Plants — Corsages
Bouquets

PAUL SCHMIDT
Florist

3121 Market St. Phone 2-4212
Youngstown, Ohio

SPENCER CORSETS
Correctly Fitted
by
HELEN MANTLE FOSTER
543 Ferndale
Phone 2-4940

POSTGRADUATE DAY — APRIL 20

Doctor!

Phone your R. direct
to us—we will deliver.

2-2483 2-1048

JONES PHARMACY
2702 Market St.

**H. H. TREUDLEY & CO.
INC.**

123 E. COMMERCE STREET
OFFICE SUPPLIES
ACCOUNT BOOKS

STEEL & WOOD OFFICE
FURNITURE

For Prompt Service
Call 3-0612

**GOLDEN AGE
GINGER ALE**

Manufactured in One of America's
Finest Beverage Plants



**GOLDEN AGE
GINGER ALE CO.**

Distributors
**Kingsbury-Schlitz-Miller's
Hi-Life and Tivoli Beers**

PHONES: 3-3333
4-4107

times in highly nervous patients a bromide quiets the cerebral activity.

Dietetic Management:

The proper use of diet in the treatment of peptic ulcer has been the subject for controversy for many years. Many clinicians have continued to employ the Sippy diet. Such diets are rarely suitable to home treatment because hospital supervision of one trained in dietetics is usually required. Many clinicians use a soft, smooth diet. The patient can eat tender meat, puree of vegetables, soft-cooked eggs, cooked fruits, milk and cream, simple puddings and gelatine. With such food there is little danger of any disturbing effect on the ulcer and in addition the diet is well balanced. As a rule it is necessary to prohibit the use of tomatoes and fresh fruit until the acute symptoms of the ulcer subside. Several disadvantages of the Sippy and similar diets lie in the fact that there is usually a deficiency in iron.

Alkali Management: There are several combinations of various alkalis used in different clinics. On these I shall elaborate later.

Methods of medical treatment:

While clinics and experienced physicians have fairly well established the mechanism by which ulcer is produced the specific cause of ulcer is still unknown. As a result treatment is in a sense symptomatic. Newer methods are constantly being advanced and there is no general agreement as to the best procedure. The doctor is therefore often at a loss as to what method of treatment to follow. Shall he continue along conventional lines or resort for example to parenteral administration of proprietary preparations now so much in vogue?

Various methods of medical treatment can be classified as follows:

1. Conventional diet—alkali—sedation (Modified Sippy regimen).
2. Administration of mucine, du-

odenal extracts, adsorbents (Kaolin and aluminum salts).

3. Duodenal or jejunal tube feedings.

4. Continuous alkalized milk or aluminum hydroxide by slow drip.

5. Parenteral administration of proteins, pepsin, insulin, vaccines, synodal, histidine hydrochloride.

6. Combined methods.

The diet alkali method of treatment is the one employed by the large majority of members of the medical profession. There is ample clinical experience and surgical evidence that any procedure which adequately will control or neutralize excess gastric acidity and secretion and which in turn will correct any impairment of gastric motor function whenever present promises the greatest success. Modification of Sippy's original method was necessary in order to avoid alkalosis and other physiological or nutritional diseases. It is also recognized that complete neutralization was not necessary to promote healing. A recent report by Emery & Monroe based on experience in 1435 cases stated that their results were satisfactory with the complete Sippy regimen in 90% of the cases, with a partial Sippy in 87% of the cases. These observers drew the conclusion that none of the present methods of treatment do more than assist in the induction of remission no matter how strict the medical schedule or how radical the operation.

The mucine method of treatment:

Mucine is a substance prepared from the mucosa of the stomach of hogs. It is useful in the treatment of peptic ulcer because it coats the ulcer and protects it against the proteolytic action of the gastric secretions and through its high combining power with free acid it unites with enough hydrochloric acid not only to neutralize the corrosive action of the gastric juices but also to prolong the rate of dialysis of pepsin through the protective layers of mucine. It is

**DON'T BE CONFUSED
BY "SIMILARITY OF NAMES"**

—it's "THE MASSACHUSETTS INDEMNITY"

There is only one company that offers to you NON-CANCEL-
LABLE and INCONTESTABLE Disability Insurance Contracts
free of restrictive limitations.

**The Massachusetts INDEMNITY Insurance
Company of BOSTON**

Phone 3-3196

•
Represented by

**LLOYD T. STILLSON, General Agent
and Associates**

**RANDY CALHOUN
JIM COZZA**

**LAMAR DONAHAY
NEIL KENNEDY**

**PETE KENNEDY
CON RYAN**

Ask Doctors who have been paid claims about our prompt efficient claim service.

POSTGRADUATE DAY — APRIL 20

• **IN ACUTE SINUSITIS**

it is often surprising the degree of relief which will sometimes follow the application of hot Antiphlogistine over the frontal and nasal regions.

Its decongestive action helps to restore drainage and relieve pain.

Antiphlogistine

**The Denver Chemical Mfg. Co.
163 Varick St., New York**

RENNER'S

**BOCK
BEER**

**HAS ALWAYS BEEN
A FAVORITE**

Order your home carton (24 bottles), in steinie or export bottles, now.

**The Renner Company
Youngstown, Ohio
Phone 44467**

April

difficult, however, to insure the maintenance of adequate amounts of mucine in the gastric contents. Only 11% of physicians to whom questionnaires had been sent noted some improvement following its use. Duodenal extract has been tried by several clinics. Two capsules of 15 grs. each were given two hours after the principal meal of the day. The results were not remarkable.

Duodenal tube feedings: It has been suggested that the most satisfactory manner of treating peptic ulcers is to pass a small tube into the duodenum so that feedings can be accomplished beyond the area of ulceration. It is assumed the ulcer will heal more readily. It has been advised by some investigators in cases in which it is difficult to control night distress. Continuous drip feeding is suggested by some of the advocates of this method of therapy. Although this method of therapy may be advisable in certain instances it is obviously impractical and unnecessary in the average case of uncomplicated peptic ulcer. 50% of physicians to whom questionnaires had been sent said that they were of the opinion that this method had some usefulness in treatment.

The continuous alkalinized milk or aluminum hydroxide method of treatment: A Rehffuss tube is passed into the stomach of the patient and a solution consisting of milk containing 5 gm. of soda bicarbonate to a quart is allowed to drip into the stomach at the rate of 30 drops a minute. Thus the patient receives three quarts of this mixture per day. Such a solution will theoretically neutralize nine quarts of 1/10 N. hydrochloric acid.

This treatment may be carried on uninterruptedly for a period of from 2-3 weeks.

The features of this type of treatment are:

1. Loss of all symptoms in from 4-6 hours after it is instituted.

2. Complete comfort of almost all patients while on the drip.

3. Eagerness to return to the drip on recurrence of even slight symptoms. In instances where no relief was obtained from the Sippy or mucine treatment immediate improvement followed the drip treatment.

The disadvantages are:

1. The method is inconvenient to the patient.

2. It is not ambulatory.

3. The tendency of the ulcer to recur after any form of therapy.

4. Probable intolerance in some patients to milk.

At the Mt. Sinai Hospital in New York 42 patients were treated by this method and they achieved results in about 70% of the cases.

Complete neutralization by the Sippy method may not always be accomplished for the following reasons:

1. Tendency of the stomach to respond to alkalosis by increased secretion of acid.

2. Secretion continues during the night in many instances and prevents healing.

3. The onset of alkalosis as a result of a large amount of alkali which may be required.

Crohn in 1929 found that colloidal aluminum hydroxide was an effective anti-acid and that it reduced total acidity to a point where complete cessation of subjective symptoms was almost an invariable rule and that it was nonabsorbable and nontoxic. Others have confirmed these statements. At Peter Bent Brigham Hospital 21 patients were treated by this method. All patients were relieved of their ulcer distress within 24 hours and had no recurrence of pain while remaining on the treatment. The only complaint of those who had followed the treatment was severe constipation in some instances. Nearly all of the patients had to take some mineral oil but clinicians believe that

NOW YOU CAN GET
**HOMOGENIZED
 VITAMIN "D" MILK**

It's easier to digest: Homogenizing breaks up the cream—by pressure—into tiny particles that form a soft curd in the stomach, making digestion easier, even for babies.

It has a new flavor: Because there is cream in every drop, this milk has a grand, new taste. Children love it and grown-ups enjoy milk more than ever.

It contains VITAMIN D: This is the precious vitamin so necessary

to develop strong bones and sound teeth. Every quart contains 400 U.S.P. units of Vitamin D.

It is more nutritious: Because it digests easily, more of the food value of this milk is assimilated by the system.

Every mother will welcome this milk that is creamy rich from top to bottom. Order Sealtest-Approved Homogenized Vitamin D Milk today.

TELEPHONE 32161



Sanitary's



Homogenized Vitamin "D" Milk

POSTGRADUATE DAY — APRIL 20



**OFFICE
 FURNITURE**

Venetian Blinds

James & Weaver

"Office Outfitters"

4-4427

W. Wood

ALCOHOLISM

—*Exclusively*—

Complete rehabilitation
 —designed to leave patient absolutely free from any craving or desire for all liquors. Desire to quit liquors our only requirement.

MAYNARD A. BUCK, M. D.

Offering Absolute Seclusion

ELM MANOR

Phone 3443

Reeves Road, Route No. 5

WARREN, OHIO

the colloidal aluminum hydroxide is a satisfactory anti-acid if given in large enough doses. At Tullane University 24 patients were treated by this method and 20 showed prompt relief of symptoms within three days.

Mutch at Royal Victoria Hospital at Montreal treated 36 patients with peptic ulcer with magnesium trisilicate and noted that following three months' observations 89% of the patients were completely free from symptoms. He stated that this alkali appears to be an efficient, non-toxic anti-acid causing rapid disappearance of symptoms, promotes healing, produces no general side effects and rarely disturbs the motility of the gastrointestinal tract.

Kraemer of the Presbyterian Hospital in Newark treated 38 private patients with the above same anti-acid. He noted improvement in all of them and believes that magnesium trisilicate is a valuable anti-acid to be used in the treatment.

The parenteral treatment:

The use of vaccines, non-specific proteins and synodal (a foreign protein) are only mentioned here as having been tried by various clinics and discarded. Larostidine, a trade name for Histidine — mono-hydrochloride — is being used by quite a number of clinicians.

Rationale for the use of Histidine as stated by some workers is that the ulcer is produced by some deficiency and that Histidine (a derivative of protein metabolism) supplies this deficiency and produces a remission of symptoms and controls the hypersecretion of acid. Others believe that Histidine stimulates an outpouring of gastric mucine which exerts a protective effect on the ulcer. At the Harper Hospital in Detroit Sandweiss reported a series of 291 cases whom he treated with the dietetic alkali management and the Histidine methods. In the former, 90% became symptom free; in the latter, 60% became symptom free. Also 22

patients were injected with distilled water and comparable results were obtained. At the Metropolitan Hospital in New York a series of cases was treated with alkali diet management and compared with the Histidine form of therapy. The Sippy form of treatment gave the majority of patients definite relief. Sterile water was used as a control and in that group slightly more so-called cures than the Histidine form of treatment were obtained. In Kansas City hospital 42 cases were treated with Larostidine. 76% obtained relief. They were of the opinion that results obtained did not justify the routine use of this procedure. At Cornell University 41 patients were treated with Histidine and they found that clinical improvement appeared to be symptomatic and transient, that it had no constant effect on the hydrochloric acid secretion and that its indications are very limited.

At the Veteran's Hospital in New Mexico Benedict treated 132 cases of peptic ulcer with Larostidine. He noted prompt improvement with ambulatory treatment and soft diet. He believes that its daily parenteral administration is the therapy of choice of peptic ulcer.

At Mayo's the treatment of peptic ulcer by Larostidine has not been satisfactory.

The combined method of treatment:

Among the many forms of medical treatment, Sippy diet, alkali regimen together with the use of sedatives still ranks highest. It has been modified to avoid nutritional disturbances and unbalance of the acid base ratio. At Massachusetts General Hospital they alternate Sippy powders with aluminum hydroxide and use the Sippy diet with sedatives. To all practical purposes this seems to be the best regimen. It is convenient, practical and produces the least number of recurrences.

This treatment adequately controls



In Our MATERNITY SHOP
Famous CAMP
Maternity Corsets
\$5

This model, with lastex groin support fitting into the side lacing, provides for extra expansion during the last period of pregnancy. Just one of many CAMP girdles for maternity and post-operative wear which our CAMP-trained corsetieres will fit to your prescription.

Maternity Shop—Second Floor

STROUSS - HIRSHBERG'S

POSTGRADUATE DAY — APRIL 20

Doctor!

Feel fully dressed—Shirts laundered at 9¢ each
 with every \$1.00 worth of dry cleaning.

THORNTON LAUNDRY AND DRY
 CLEANING COMPANY

Phone 40155

Girard 139

**ALWAYS
 DEPENDABLE
 PRODUCTS**

Prescribe or Dispense ZEMMER
 Pharmaceuticals. . . Tablets, Lozenges,
 Ampoules, Capsules, Ointments, etc.
 Guaranteed reliable potency. Our
 products are laboratory controlled.

Write for catalog.

Chemists to the Medical Profession.

THE ZEMMER COMPANY, Oakland Station, PITTSBURGH, PA.

symptoms of distress and complications in the acute stage of ulcers. It insures adequate healing by the application of physiological principles such as neutralization of acid and it decreases the peptic activity. Finally, together with sane living, adequate rest, supplementary feedings, avoidance of infection and non-stimulating diet, it prevents recurrences.

The treatment of peptic ulcer with hemorrhage:

At the Crile Clinic patients are put to bed at once, morphine and atrophine are given and the patient is starved for from 24-48 hours. If bleeding will not cease a careful gastric lavage with 4-6 ounces of ice cold water is tried. The wash fluid may consist of 1-1000 ferric chloride followed by the installation of 1 dram of 1-1000 of adrenalin. Blood transfusions are indicated if the hemoglobin falls to 40-50%.

Neulengrascht of Copenhagen treated 251 patients with bleeding duodenal ulcers by giving the patient a full puree diet with soda bicarbonate and magnesium subcarbonate and extract of hyoscyamine and ferric lactate. They were allowed to eat as much as they wanted. A few patients were given blood transfusions on admission. He had a mortality of 1%. Christianson of the same city treated 289 cases by the starvation method and had a mortality of 8%. Neulengrascht believes that many patients with bleeding ulcer die of exhaustion and not of hemorrhage. Patients past 50 years of age who have massive hemorrhages should be operated upon as soon as possible.

In the past five years 163 cases of peptic ulcer were treated at St. Elizabeth's Hospital. 55% of these were confirmed by x-ray. In this series of cases 79% were duodenal ulcer and 21% were gastric ulcer. There were 26 cases of peptic ulcer with hemorrhage with two deaths; a mortality of 7½%. In 80% of these cases the

treatment was sedation, transfusions and Sippy diet. The rest of the cases were treated by transfusion, starvation and sedation. There were 35 perforations of peptic ulcer with ten deaths; a mortality of 29%.

Of the peptic ulcers treated medically 20% were treated with Creamalin and modified Sippy diet and antispasmodics, 60% with Sippy powders and Sippy diet, 17% with soft diet alone and 3% with Larostidine and soft diet. In all the cases the charts recorded improvement. We were unable to check the follow-up results. These are just to show the type of treatment that has been instituted in this hospital over the past five years.

In closing I would like to enumerate some of the measures which should govern and aid in the treatment of peptic ulcer, namely:

1. Individualize your case.
2. Treat the patient together with the ulcer.
3. Inasmuch as it is economically possible try to relieve the patient's mind from anxiety and worry.
4. Change the surroundings.
5. Demand the patient's cooperation, if not he will soon see another doctor.
6. Hospitalize acute cases.
7. Use alkalis, sedatives and antispasmodics to control symptoms. Be on guard for alkalosis.
8. A vacation may be your best alkali and antispasmodic available.
9. Use Sippy or a modified Sippy diet early and gradually work into a soft, bland diet.
10. Permit no smoking or alcohol.
11. Be on the look-out for complications such as hemorrhage and perforation and when found do not temporize but call in a surgical consultant.
12. Follow-up your patient as often as you think necessary for recurrences frequently occur. If possible check with x-rays. Finally:
13. Give your patient a certain amount of confidence in his own ability to get well.

NEWS

Dr. and Mrs. R. E. Odom are back from a vacation at Palm Beach, Florida.

Among those vacationing in Florida are Dr. and Mrs. J. B. Kupec.

Dr. E. C. Rinehart had the misfortune to fall and break his leg and is confined to the South Side Hospital.

Dr. J. P. Harvey is doing nicely after an appendectomy performed at the North Side a few days ago.

Dr. John Noll is attending a meeting of the American College of Physicians, New Orleans.

Dr. and Mrs. Colin R. Clark are home after spending a vacation at Atlantic City.

Dr. John M. McCann attended the American College of Physicians meeting at New Orleans.

Drs. Turner and McNamara spent a few days in Indianapolis attending a meeting of the American College of Surgeons.

Dr. Claude B. Norris addressed the Marion County Medical Society, April 4th, his subject being, "Fungus Infection of the Skin." Other addresses by Dr. Norris are:

April 6th, the Youngstown Technicians Society, subject: "Important Technical Procedure, Useful in Dermatology."

April 11th, the Salem and Sebring Rotary Clubs, subject: "Syphilis."

April 14th, Dr. Norris will address the Ashland County Medical Society, his subject being "The Management of Common Skin Disorders."

**YOU WRITE IT—
WE FILL IT CORRECTLY.**

Phone 27744

DOBSON'S PHARMACY

100 E. Midlothian

We Deliver

Drink Pepsi Cola

Bottled & Distributed by

The J. F. Giering Bottling Co.

Youngstown, Ohio

Phone 6-2212

**This publication is printed by
The Youngstown Printing Co.**



**The
Youngstown Arc Engraving Co.**
Vindicator Square
Youngstown, Ohio

SORICIN



IN INTESTINAL TOXEMIA

Proved in 17,862 Clinical Cases

A TABULATION of reports from fifteen clinical investigators covers 17,862 cases in which Soricin was used in treatment of migraine, mucous colitis, urticaria, vertigo, gastrointestinal allergy, functional diarrhea, irritable colon, and similar conditions caused by intestinal absorption of toxic substances.

Complete relief or improvement was obtained in 80% to 90% of cases treated.

Soricin acts by reducing fecal toxicity and preventing absorption of toxic matter from the bowel. Soricin is a non-toxic castor oil soap, and does not interfere

with normal gastrointestinal digestion.

Why not try Soricin for that frequently encountered patient who complains of headache, vertigo, nervousness, lassitude, and flatulence, usually associated with vague abdominal discomfort? There may be one in your waiting room now.

Soricin (brand of sodium ricinoleate) is available in 5 grain enteric coated tablets. Other dosage forms include Soricin and Bile Salts Tablets and Soricin, Bile Salts and Pancreatin Tablets. All are supplied in bottles of 100. Average dosage: four to eight tablets daily as indicated.

Write for clinical tabulation, sample, and complete literature

The Wm. S. Merrell Company

Founded 1828 • Cincinnati, U. S. A.

There is a Council-
Accepted high potency
fish liver oil available
that is advertised only
to the medical profes-
sion and not exploited
to the laity... It is called

OLEUM PERCOMORPHUM

(Liquid and Capsules)

Specify MEAD'S



Yours for Keeping the Faith

MEAD JOHNSON & COMPANY

EVANSVILLE, INDIANA, U.S.A.